

SINGAPORE GUIDELINES ON SEXUAL HEALTH FOR WOMEN OF REPRODUCTIVE AGE

The College of Obstetricians & Gynaecologists Singapore (COGS) Sexual Health Guidelines for Women of Reproductive Age in Singapore Work Group:

Tse Yeun TAN‡, Jean-Jasmin Mi-Li LEE‡, Caroline CHUA, Xinyi CHEONG, Lay Kok TAN, Jason LIM, Suraj KUMAR, Zhi Ying CHIE, Jay LIM, Benny Kai Guo LOO, Weiliang CHONG, Wan Hui YIP, Melissa CHIANG, Delphine KOH, Zhong Wei HUANG, June ENG, Chee Wai KU, Phaik Ling Elaine QUAH, Kok Hian TAN

‡ These authors contributed equally as first authors.

INTRODUCTION

Sexual health is a fundamental aspect of overall health and well-being. According to the World Health Organization (WHO)¹, sexual health is defined as "a state of physical, emotional, mental, and social well-being related to sexuality; it is not merely the absence of disease, dysfunction, or infirmity." Achieving sexual health requires a positive and respectful approach to sexuality and sexual relationships, as well as the possibility of pleasurable and safe sexual experiences, free from coercion, discrimination, and violence. For sexual health to be attained and maintained, the sexual rights of all individuals must be respected, protected, and fulfilled.

Healthy sexual expression of sexuality offers inherent physical and mental health benefits, contributing to emotional intimacy, relationship satisfaction, and overall quality of life². Female sexual function is also a vital component of this holistic understanding of sexual health. It plays a significant role in fertility and reproductive health, which is especially pertinent when trying to conceive, during pregnancy, and after childbirth³⁻⁵.

Despite its importance, sexual dysfunction in women is often underrecognized and undertreated. Studies worldwide report varying prevalence rates of sexual dysfunction among women of reproductive age, with figures ranging from 27% to over 50%^{4,5}. Common issues include low sexual desire, arousal difficulties, orgasmic disorders, and sexual pain^{4,5}. The variability in definitions and diagnostic criteria for sexual concerns, difficulties, disorders, and dysfunctions (SCDDD) contributes to the challenges in identifying and addressing these issues effectively⁴.

Many women do not perceive their sexual health concerns as medical issues or may feel hesitant to discuss them with healthcare professionals (HCPs)⁶. Barriers include uncertainty about initiating the conversation, confusion about which clinician to consult, and societal stigmas surrounding sexual topics^{7,8}. Moreover, HCPs may avoid addressing sexual health due to personal discomfort, lack of training, time constraints, or concerns about professional boundaries⁷⁻⁹.

Addressing female sexual health is essential for improving overall health, relationships, and reproductive outcomes¹⁰. HCPs are well-positioned to enhance the quality of care for women by integrating sexual health into routine practice^{10,11}. This requires a multidisciplinary approach and should be considered a legitimate and important aspect of clinical care throughout a woman's reproductive life^{12,13}.

DEFINITIONS AND CLASSIFICATION

Female Sexual Dysfunction (FSD) encompasses various disorders related to desire, arousal, orgasm, and sexual pain. According to the Diagnostic and Statistical Manual of Mental Disorders, Fifth Edition (DSM-5)¹⁴, FSD includes:

- **Female Sexual Interest/Arousal Disorder (FSIAD):** A lack or significant reduction in sexual interest/arousal.
- **Female Orgasmic Disorder:** Difficulty in achieving orgasm or reduced orgasmic intensity.
- **Genito-Pelvic Pain/Penetration Disorder (GPPPD):** Pain during intercourse or penetration attempts, fear or anxiety about pain, or tensing of pelvic floor muscles.

For a diagnosis of FSD, symptoms must cause significant distress, occur at least 75% of sexual events, and persist for a minimum of six months.

These guidelines aim to provide HCPs with an overview of sexual health issues in reproductive-age women and to help them develop the knowledge and comfort necessary to identify and manage women with sexual concerns. The goal is to offer guidance on screening, education, management, and referral for women experiencing sexual problems.

TARGET AUDIENCE

These guidelines are intended for all HCPs who interact with reproductive-age women in their daily practice, including general practitioners, family physicians, obstetricians and gynaecologists, nurses, midwives, pelvic health physiotherapists, and mental health professionals such as psychiatrists, psychologists, counsellors, and social workers.

GUIDELINES OVERVIEW

The guidelines are divided into the following sections:

1. Screening
2. Assessment
3. Education
4. Optimization
5. Preconception Considerations
6. Antenatal Considerations
7. Postnatal Considerations
8. Special Considerations

1. Screening

Recognizing the importance of sexual health is the first step in addressing sexual concerns among reproductive-age women. Sexual health is integral to overall health and is closely intertwined with physical, emotional, and mental well-being^{1,2,4}. It encompasses not only medical topics such as contraception and sexually transmitted infections but also includes sexual functioning, which involves desire, arousal, orgasm, and satisfaction^{3,4}.

Women should be made to feel comfortable as much as possible when discussing sexual health, including sexual functioning, with HCPs^{4,15}. Many women expect HCPs to initiate discussions when appropriate about sexual health⁹. Therefore, it is important for HCPs to proactively engage patients in conversations about their sexual well-being^{4,9,15}. Incorporating questions about sexual activity, contraceptive use, sexually transmitted infections, pregnancy planning, sexual function, and sexual concerns into routine history-taking can facilitate these discussions when appropriate within the clinical context^{4,7}.

HCPs who may be consulted by women regarding sexual health include doctors, particularly gynaecologists and primary care providers such as general practitioners and family physicians^{4,7,13,16}. Nurses, midwives, physiotherapists specializing in women's and pelvic health, psychiatrists, psychologists, and counsellors also play significant roles in addressing women's sexual health concerns^{7,12,13}.

It is essential for HCPs to regard the identification and management of a woman's sexual health issues as important and legitimate elements of her clinical care^{7,16}. By normalizing discussions about sexual health and demonstrating openness and sensitivity, HCPs can create a safe environment for women to express their concerns¹⁷.

2. Assessment

Assessment of female sexual dysfunction can be undertaken using brief questionnaires such as the Female Sexual Function Index-6 (FSFI-6)¹⁸. These tools help to identify specific areas of concern and facilitate more targeted interventions.

Involving the woman's partner in addressing sexual concerns can be beneficial when it is appropriate and safe to do so^{4,19}. Male sexual issues can also affect the female partner's sexual function and vice versa^{4,19}. Communication within the couple has been linked to various aspects of sexual functioning, including sexual desire, arousal, lubrication, orgasm, satisfaction, erectile function, and the experience of pain²⁰.

The **PLISSIT** (Permission, Limited Information, Specific Suggestions, Intensive Therapy) model²¹ can be used for preliminary screening and intervention. This approach starts with giving the patient permission to express sexual symptoms or concerns, offering information about how certain changes can affect sexuality or sexual functioning, providing specific suggestions to address these issues, and referring for intensive therapy if necessary²¹.

HCPs should distinguish between sexual concerns or difficulties and sexual dysfunction. Many sexual problems are transient and may resolve with basic education or minor interventions²².

Early and appropriate intervention may allow resolution of sexual concerns and prevent further progression to sexual dysfunction²².

It is also important to consider the implications of medical conditions and their treatments on women's sexuality⁴. Chronic illnesses such as hypertension, coronary heart disease, diabetes, and thyroid dysfunction can cause physical and emotional changes that affect sexual functioning^{4,23}. Clinicians caring for women with chronic illnesses should integrate information about sexual care into their medical therapy or offer a referral to specialist sexual health services when appropriate²³.

3. Education

HCPs should encourage women to use condoms consistently and to take other steps to promote sexual health and prevent sexually transmitted infections, such as receiving the human papillomavirus (HPV) vaccination if they are sexually active^{4,23}. It is important to educate patients about normal fluctuations in sexual function response due to factors such as the menstrual cycle, stress, or relationship dynamics^{4,23}. For instance, patients should be informed that it is common to experience changes in sexual function related to the menstrual cycle, periods of stress, or interpersonal conflict⁴. Sexual frequency may also change with age and the duration of the relationship²³.

Changes in sexual functioning should be treated only if the woman expresses distress about these changes and desires treatment^{4,23}. HCPs should offer information about the availability and efficacy of treatments and respect the patient's preferences and autonomy in decision-making.

4. Optimization

A holistic approach to optimizing sexual health involves addressing medical, psychological, sociocultural and relational factors that can affect sexual function.

Medical conditions such as hypertension, coronary heart disease, diabetes, obesity, and thyroid dysfunction can affect sexual functioning^{4,24}. HCPs should be aware of these implications and integrate information about sexual care into their treatment plans^{23,24}. Optimizing the management of these conditions can improve overall well-being and sexual function.

Gynaecological conditions like endometriosis and adenomyosis can also affect sexual functioning²⁵. Early diagnosis and multidisciplinary management can prevent impairment of sexual quality of life^{11,13}. Symptoms such as menstrual irregularities, pelvic pain, dyspareunia, dysmenorrhea, and subfertility are common²⁶. Encouraging patients to communicate sexual problems as part of the assessment and referring them to a gynaecologist for further evaluation and treatment is important.

HCPs should inquire about the presence of any sexual dysfunction in male partners, when possible, as male sexual issues can affect the female partner's sexual function^{4,19}. Providing resources for andrology services can help address these concerns.

HCPs should recommend regular physical exercise²⁷⁻³⁰ and appropriate pelvic floor muscles training³¹⁻³⁵ to enhance overall well-being and sexual function. Referrals to pelvic health physiotherapy should be provided as needed and may be appropriate for co-managing pelvic floor disorders such as overactive bladder, urinary incontinence, prolapse, and pelvic pain³⁵⁻³⁷.

Sexual concerns that do not respond to first-line interventions or cases of sexual dysfunction should be referred to HCPs who specialize in sexual health³⁸. Establishing a list of clinical sexual health resources in the community can facilitate timely referrals when necessary.

Oral contraceptives can cause sexual dysfunction in reproductive-age women. HCPs should inquire about sexual function before prescribing oral contraceptives and assess sexual function at initiation and at regular intervals thereafter³⁹. This approach allows contraceptive-related sexual dysfunction to be managed early, such as by switching the patient to newer-generation contraceptives or other forms of contraception⁴⁰.

Maintaining good mental health is associated with optimal sexual functioning⁴¹. The experience of psychosocial and interpersonal factors, such as daily life stressors, caregiver duties, family responsibilities, or major life events, can negatively impact sexual functioning⁴¹. HCPs should explicitly ask about the experience of stressors⁴² or use brief screeners such as the Depression Anxiety Stress Scales (DASS-10)⁴³ to screen for chronic or prolonged stress. Referral to counselling or psychology services should be offered as desired.

Relationship factors like poor communication about sexual issues, conflict, and dissatisfaction can negatively influence sexual functioning⁴⁴. Addressing these issues and offering referrals to couples counselling or psychology services can be beneficial.

Mental disorders (e.g., depressive or bipolar disorder, anxiety disorder, posttraumatic stress disorder, psychotic disorder) along with their treatments, can affect sexual functioning⁴⁵. HCPs should inquire about the presence of any mental health condition and obtain specific information about psychotropic treatments⁴⁵. Informing patients of the link between mental health conditions, their treatment, and sexual dysfunction, and offering referrals to specialist sexual health services for further management, is important.

In cases where poor sexual functioning is linked to past or ongoing sexual or physical trauma, the healthcare provider should consider a referral to agencies that specialize in trauma-focused assessment or support¹⁷.

5. Preconception Considerations

When trying to conceive, sexual concerns can increase due to heightened stress⁴⁶. HCPs should ask about stress related to the process of trying to conceive or use brief screeners like the DASS-10⁴³ to assess for chronic stress. Infertility, defined as the failure to achieve

pregnancy after 12 months or more of regular unprotected intercourse (or six months for women aged 35 and older), can cause distress and affect sexual functioning²⁵. Early evaluation and treatment may be warranted based on medical history and age.

In assessing sexual health concerns, HCPs should involve the woman's partner when appropriate. Administering questionnaires like the FSFI-6¹⁸, can provide more information about sexual concerns.

Preconception counselling on sexual function is important⁴⁷. Couples trying to conceive are encouraged to have regular intercourse, and reproductive efficiency is highest with intercourse every one to two days during the fertile window⁴⁷. However, the optimal frequency of intercourse is best defined by the couple's preference within that context⁴⁷.

Patients should be informed that coital position does not affect fecundability, and remaining supine after intercourse does not facilitate sperm transport⁴⁷. Although some lubricants may affect sperm parameters in vitro, their use does not significantly impact fertility in practice⁴⁷. HCPs should inform patients that sexual functioning may be affected as routines become ritualized and stress increases during efforts to conceive⁴⁷. Support should be offered if desired⁴⁷.

Optimizing preconception sexual health involves addressing medical conditions that can affect sexual function, such as chronic illnesses and their treatments^{24,27-29}. Inquiring about male partner's sexual dysfunction and providing resources for andrology services can be helpful^{4,19}.

Maintaining good mental health is crucial⁴². HCPs should ask about psychosocial and interpersonal factors that may impact sexual functioning, such as stress and relationship issues^{41,42,44,45}. Addressing mental health conditions and offering referrals to counselling or psychology services is important^{4,19}.

6. Antenatal Considerations

Pregnancy introduces biological, psychological, and social changes that may alter sexual health^{48,49}. Sexual function often declines during pregnancy⁴⁹ and may not return to baseline levels during the postpartum period⁴⁸. Factors such as hyperemesis in the first trimester, fear of causing obstetric complications, physical changes, difficulty with coital positioning, changes in body image, fatigue, and urinary complaints contribute to this decline⁴⁸.

HCPs should screen for sexual functioning during antenatal visits in any trimester. Questions about sexual well-being should be part of routine medical health questioning. Using the PLISSIT²¹ approach can facilitate discussions.

Proactively informing patients and their partners that HCPs are available to address concerns about sex during pregnancy is important^{50,51}. Sexual functioning should be discussed early in prenatal care, before hospital discharge postpartum, and at postnatal check-ups^{50,51}. Patients should be educated that their baseline for comparison is sexual activity before pregnancy^{50,51}.

If there are no medical or obstetric complications, couples can be reassured that they can continue sexual activity if they desire⁵². Education about common changes in sexual function and frequency across pregnancy trimesters should be provided⁵¹. Advice on adapting coital positions and using lubricants to address dyspareunia can support sexual adjustment⁵¹.

Assessment for sexual functioning can be conducted with questionnaires like the FSFI¹⁸. Including the partner's sexual functioning in the assessment can provide a comprehensive understanding^{4, 19}. Diagnosing sexual dysfunction is important if difficulties have persisted for six months.

Addressing issues like vulvar pain involves advising women to avoid irritants and providing symptomatic treatment⁵³. Referral to pelvic health physiotherapy can prepare women with GPPPD for normal vaginal delivery⁵³. Recommending the use of lubricants or vaginal moisturizers for vaginal dryness can alleviate discomfort^{4,52}.

Treating medical, psychological, and relationship problems, and addressing sociocultural issues can help women and their partners deal with low desire and orgasm difficulties during pregnancy⁵⁴. Encouraging exercise⁵⁵ and pelvic floor strengthening exercises^{33,56} can improve well-being and maintain sexual function during pregnancy⁵⁷.

HCPs should screen for antenatal depression and anxiety using tools like the Edinburgh Postnatal Depression Scale⁵⁸. Referrals to mental health professionals should be made when necessary.

7. Postnatal Considerations

Postpartum women often experience significant changes in sexual function, including dyspareunia, lack of lubrication, difficulty reaching orgasm, vaginal bleeding, and loss of desire⁸. Sexual dysfunction prevalence rates are high in the postpartum period and can significantly impact quality of life⁸.

Screening for sexual functioning should occur during postpartum medical reviews. Questions about resumption of sexual activity and sexual difficulties should be incorporated into routine health questioning, using the PLISSIT²¹ approach to facilitate discussions.

Assessment can include using the FSFI¹⁸ to ascertain sexual dysfunction. Ensuring that perineal injuries have healed and addressing any bladder or bowel difficulties is essential⁵⁸. Referral to specialists may be necessary for managing complications.

Education about resuming sexual intercourse, typically recommended at least four to six weeks postpartum when healing is complete, is important.⁵⁹ Explaining how hormonal changes, breastfeeding, contraception, and fatigue can affect sexual functioning helps women understand and manage these changes⁶⁰.

Optimizing sexual health involves referring women to pelvic health physiotherapy for management of post-delivery trauma and pelvic floor rehabilitation^{61,62}. Addressing lifestyle

factors, such as stress and fatigue, can improve sexual desire and arousal^{59,63}. Advising women to make time for themselves can promote well-being^{59,63}.

Holistic management includes recommending gradual exercise introduction based on postpartum physical activity guidelines⁶⁴ and screening for postpartum depression using tools like the Edinburgh Postnatal Depression Scale⁵⁷. Addressing mental health is crucial, as postpartum depression can present with sexual difficulties⁶⁰.

8. Special Considerations

Women who have a history of abuse, including physical, emotional, or sexual abuse are more likely to experience sexual dysfunction^{17,65,66}. It's important to consider the impact of adverse events, such as signs of post-traumatic stress disorder (PTSD), as PTSD is associated with poorer sexual functioning^{17,65,66}. Referrals to agencies that provide specialized trauma-focused support should be considered¹⁷.

In cases where patients present with high-risk concerns such as ongoing suicidal ideations, referrals to psychiatrists for risk management are necessary.

HCPs should be inclusive and sensitive to the needs of Lesbian, Gay, Bisexual, Trans, Queer or Questioning+ (LGBTQ+) individuals, by acknowledging that they may face unique sexual health concerns⁶⁷. Providing appropriate resources and referrals is important⁶⁷.

Women who have undergone cancer treatments may face specific sexual health challenges due to the effects of chemotherapy, radiation, or surgery on hormonal levels, physical function, and emotional well-being.⁶⁸ Addressing these challenges and providing referrals for specialized support is essential.⁶⁸

There are women who have other unique sexual health concerns associated with specific physical, cognitive and/or medical challenges. It is important to recognise these concerns and provide referrals for specialized support as needed.⁶⁹

CONCLUSION

Female sexual health is a critical aspect of women's overall well-being and quality of life¹⁻³. HCPs have a vital role in identifying, addressing, and managing sexual health concerns^{10,11}. By integrating sexual health into routine clinical practice and adopting a patient-centred approach, HCPs can enhance the care provided to reproductive-age women.

These guidelines serve as a comprehensive resource to support HCPs in delivering effective sexual healthcare. Ongoing education, interdisciplinary collaboration, and the establishment of referral networks are essential for providing holistic care.

REFERENCES

1. Defining sexual health [Internet]. Who.int. [cited 2024 Dec 2]. Available from: <https://www.who.int/teams/sexual-and-reproductive-health-and-research/key-areas-of-work/sexual-health/defining-sexual-health>
2. Dehghankar L, Panahi R, javanmardi3 E, Khatooni M, Valinezhad S, Anbari M. The effect of women's sexual functioning on quality of their sexual life [Internet]. 2021; Available from: <http://dx.doi.org/10.15167/2421-4248/JPMH2021.62.3.1945>
3. Gianotten WL, Alley JC, Diamond LM. The health benefits of sexual expression. Int J Sex Health [Internet]. 2021;33(4):478–93. Available from: <http://dx.doi.org/10.1080/19317611.2021.1966564>
4. Wylie K. Assessment & management of sexual problems in women. Journal of the Royal Society of Medicine [Internet]. 2007 Dec;100(12):547–50. Available from: <http://dx.doi.org/10.1177/0141076807100012011>
5. Alidost F, Pakzad R, Dolatian M, Abdi F. Sexual dysfunction among women of reproductive age: A systematic review and meta-analysis. International Journal of Reproductive BioMedicine (IJRM) [Internet]. 2021 Jun 23 Available from: <http://dx.doi.org/10.18502/ijrm.v19i5.9251>
6. Wendt E, Hildingh C, Lidell E, Westerståhl A, Baigi A, Marklund B. Young women's sexual health and their views on dialogue with health professionals. Acta Obstet Gynecol Scand [Internet]. 2007;86(5):590–5. Available from: <http://dx.doi.org/10.1080/00016340701214035>
7. Kingsberg SA, Schaffir J, Faught BM, Pinkerton JV, Parish SJ, Iglesia CB, et al. Female sexual health: Barriers to optimal outcomes and a roadmap for improved patient–clinician communications. J Womens Health (Larchmt) [Internet]. 2019;28(4):432–43. Available from: <http://dx.doi.org/10.1089/jwh.2018.7352>
8. Barrett G, Pendry E, Peacock J, Victor C, Thakar R, Manyonda I. Women's sexual health after childbirth. BJOG: An International Journal of Obstetrics and Gynaecology. 2000 Feb;107(2):186–95.
9. Haboubi NHJ, Lincoln N. Views of health professionals on discussing sexual issues with patients. Disability and Rehabilitation [Internet]. 2003 Jan;25(6):291–6. Available from: <http://dx.doi.org/10.1080/0963828021000031188>
10. World Health Organization. Brief sexuality-related communication : recommendations for a public health approach. Geneva, Switzerland: World Health Organization; 2015. Available from: <https://ddei5-0-ctp.trendmicro.com:443/wis/clicktime/v1/query?url=https%3a%2f%2fwww.ncbi.nlm.nih.gov%2fbooks%2fNBK311026%2f&umid=CCB8807A-22ED-B506-AB07-DB803735E8B3&auth=506d7fb60e41bcf4ca9a42eb3a951ef6efc5338b-9916784539c773fb532e94b39f5cab806e009dfd>
11. Berman L, Berman J, Felder S, Pollets D, Chhabra S, Miles M, et al. Seeking help for sexual function complaints: what gynecologists need to know about the female patient's experience. Fertility and Sterility. 2003 Mar;79(3):572–6.
12. Davis S. Rehabilitation: the use of theories and models in practice. Edinburgh: Elsevier Churchill Livingstone; 2006.
13. Nusbaum MR, Gamble G, Skinner B, Heiman J. The high prevalence of sexual concerns among women seeking routine gynecological care. PubMed. 2000 Mar 1;49(3):229–32.

14. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th-TR. American Psychiatric Association; 2022.
15. Kingsberg SA. Taking a Sexual History. *Obstetrics and Gynecology Clinics of North America*. 2006 Dec;33(4):535–47.
16. Stanley EE, Pope RJ. Characteristics of Female Sexual Health Programs and Providers in the United States. *Sexual Medicine*. 2022 Aug;10(4):100524.
17. Lafrenaye-Dugas AJ, Hébert M, Godbout N. Sexual satisfaction improvement in patients seeking sex therapy: evaluative study of the influence of traumas, attachment and therapeutic alliance. *Sexual and Relationship Therapy*. 2020 Feb 12;1–18.
18. Isidori AM, Pozza C, Esposito K, Giugliano D, Morano S, Vignozzi L, et al. Development and Validation of a 6-Item Version of the Female Sexual Function Index (FSFI) as a Diagnostic Tool for Female Sexual Dysfunction. *The Journal of Sexual Medicine*. 2010 Mar;7(3):1139–46.
19. Yeoh SH, Razali R, Sidi H, Razi ZRM, Midin M, Jaafar NRN, et al. The relationship between sexual functioning among couples undergoing infertility treatment: a pair of perfect gloves. *Comprehensive Psychiatry*. 2014 Jan;55:S1–6.
20. Mallory AB, Stanton AM, Handy AB. Couples' sexual communication and dimensions of sexual function: A meta-analysis. *J Sex Res* [Internet]. 2019;56(7):882–98. Available from: <http://dx.doi.org/10.1080/00224499.2019.1568375>
21. Annon JS. The PLISSIT model: A proposed conceptual scheme for the behavioral treatment of sexual problems. *J Sex Educ Ther* [Internet]. 1976;2(1):1–15. Available from: <http://dx.doi.org/10.1080/01614576.1976.11074483>
22. Mercer CH, Fenton KA, Johnson AM, Wellings K, Macdowall W, McManus S, Nanchahal K, Erens B. Sexual function problems and help seeking behaviour in Britain: national probability sample survey. *Bmj*. 2003 Aug 21;327(7412):426-7.
23. Lamont J, Bajzak K, Bouchard C, Burnett M, Byers S, Cohen T, Fisher W, Holzapfel S, Senikas V. Female sexual health consensus clinical guidelines. *Journal of obstetrics and gynaecology Canada* [Internet]. 2012 Aug 1;34(8):769-75. Available from: [http://dx.doi.org/10.1016/s1701-2163\(16\)35341-5](http://dx.doi.org/10.1016/s1701-2163(16)35341-5)
24. Das J, Yadav S, Arora B. Comorbidities of Female Patients with Sexual Dysfunction in a Psychiatry Clinic: A Cross-Sectional Study. *Journal of Psychosexual Health*. 2022 May 22;4(3):162–70.
25. Pluchino N, Wenger JM, Petignat P, Tal R, Bolmont M, Taylor HS, Bianchi-Demicheli F. Sexual function in endometriosis patients and their partners: effect of the disease and consequences of treatment. *Human reproduction update*. 2016 Nov 20;22(6):762-74.
26. Lara LA. Sexuality in couples with reproductive difficulties. *Female Sexual Function and Dysfunction*. 2017:107-22.
27. Greco A, Zentner A, Brotto LA. Comparison of Medical Management versus Bariatric Surgery for Obesity Management: Effects on Sexual Function. *Journal of Sex & Marital Therapy*. 2021 Jun 26;47(7):721–30.
28. Mosallanezhad Z, Honarmand F, Poornowrooz N, Jamali S. The relationship between body mass index, sexual function and quality of life in women of reproductive age in Iran. *Sexual and Relationship Therapy*. 2020 Feb 9;1–11.
29. Gao Z, Liang Y, Deng W, Qiu P, Li M, Zhou Z. Impact of Bariatric Surgery on Female Sexual Function in Obese Patients: a Meta-Analysis. *Obesity Surgery*. 2019 Oct 29;30(1):352–64.

30. Vanti C, Ferrari S, Chiodini M, Cesare Olivoni, Bortolami A, Paolo Pillastrini. Sexual Disability in Low Back Pain: Diagnostic and Therapeutic Framework for Physical Therapists. *Healthcare* [Internet]. 2023 Dec 29 [cited 2024 Jun 25];12(1):80–0. Available from: <https://doi.org/10.3390%2Fhealthcare12010080>
31. Tabatabaie NS, Borujeni FR, Tavakol Z, Shahraki HR. The Effect of Yoga and Pelvic Floor Muscle Exercise on Sexual Function and Sexual Self-esteem of Reproductive-age Women: A Randomized Clinical Trial. *International Journal of Community Based Nursing and Midwifery*. 2024 Jul;12(3):188.
32. Basgol S, Oskay U. Examining the effectiveness of home-based pelvic floor muscle training in treating sexual dysfunction in women. *Int J Caring Sci*. 2016 Jan 1;9(1):135-43.
33. Albaugh J. Female sexual dysfunction. *International Journal of Urological Nursing*. 2014 Mar;8(1):38-43.
34. Nygaard AS, Rydningen MB, Stedenfeldt M, Wojniusz S, Larsen M, Lindsetmo R, et al. Group-based multimodal physical therapy in women with chronic pelvic pain: A randomized controlled trial. *Acta Obstetricia et Gynecologica Scandinavica*. 2020 Jun;99(10):1320–9.
35. Trahan J, Leger E, Allen M, Koebele R, Yoffe MB, Simon C, et al. The Efficacy of Manual Therapy for Treatment of Dyspareunia in Females. *Journal of Women's Health Physical Therapy*. 2019;43(1):28–35.
36. Hill DA, Taylor CA. Dyspareunia in women. *American family physician*. 2021 May 15;103(10):597-604.
37. Stein A, Sauder SK, Reale J. The Role of Physical Therapy in Sexual Health in Men and Women: Evaluation and Treatment. *Sexual Medicine Reviews*. 2019 Jan;7(1):46–56.
38. Kingsberg SA, Adler B, Metropoulos J, Faubion SS. The yin and yang of GSM and low sexual desire. *Climacteric*. 2023 Apr 21;26(4):323–8.
39. Lee JJ, Low LL, Ang SB. Oral contraception and female sexual dysfunction in reproductive women. *Sexual Medicine Reviews*. 2017 Jan;5(1):31-44.
40. Lee J, Tan T, Ang S. Female sexual dysfunction with combined oral contraceptive use. *Singapore Medical Journal*. 2017 Jun;58(6):285–8.
41. Kalmbach DA, Pillai V. Daily affect and female sexual function. *The journal of sexual medicine*. 2014 Dec;11(12):2938-54.
42. Hamilton LD, Meston CM. Chronic stress and sexual function in women. *The journal of sexual medicine*. 2013 Oct;10(10):2443-54.
43. Kim Halford W, Frost ADJ. Depression Anxiety Stress Scale-10: A Brief Measure for Routine Psychotherapy Outcome and Progress Assessment. *Behaviour Change*. 2021 Aug 11;38(4):1–14.
44. McCabe MP, Connaughton C. Sexual dysfunction and relationship stress: how does this association vary for men and women?. *Current opinion in psychology*. 2017 Feb 1;13:81-4.
45. Montejo AL, Montejo L, Baldwin DS. The impact of severe mental disorders and psychotropic medications on sexual health and its implications for clinical management. *World psychiatry*. 2018 Feb;17(1):3-11.
46. Loy SL, Ku CW, Cheung YB, Godfrey KM, Chong YS, Shek LPC, et al. Fecundability in reproductive aged women at risk of sexual dysfunction and associated risk factors: a prospective preconception cohort study. *BMC Pregnancy and Childbirth*. 2021 Jun 25;21(1).

47. Practice Committee of the American Society for Reproductive Medicine. Optimizing natural fertility: a committee opinion. *Fertility and Sterility*. 2022 Jan 1;117(1):53-63.
48. Banaei M, Azizi M, Moridi A, Dashti S, Yabandeh AP, Roozbeh N. Sexual dysfunction and related factors in pregnancy and postpartum: a systematic review and meta-analysis protocol. *Systematic reviews*. 2019 Dec;8:1-5.
49. Nezal AJ, Rad FS, Kalhor M, Alipour M, Montazeri A. Sexual quality of life in pregnant women: A cross sectional study. *Payesh (Health Monitor)*. 2018 Aug 15;17(4):421-9.
50. Staruch M, Kucharczyk A, Zawadzka K, Wielgos M, Szymusik I. Sexual activity during pregnancy. *Neuroendocrinol Lett*. 2016 Jan 1;37(1):53-8.
51. Geuens S, Polona Mivšek A, Gianotten W. *Midwifery and Sexuality*. Springer Nature; 2023.
52. MacPhedran SE. Sexual activity recommendations in high-risk pregnancies: What is the evidence?. *Sexual medicine reviews*. 2018 Jul;6(3):343-57.
53. Dias-Amaral A, Marques-Pinto A. Female Genito-Pelvic Pain/Penetration Disorder: Review of the Related Factors and Overall Approach. *Revista Brasileira de Ginecologia e Obstetrícia / RBGO Gynecology and Obstetrics* [Internet]. 2018 Nov 14 [cited 2019 Nov 2];40(12):787–93. Available from: <http://www.scielo.br/pdf/rbgo/v40n12/0100-7203-rbgo-40-12-00787.pdf>
54. Daescu AM, Navolan DB, Dehelean L, Frandes M, Gaitoane AI, Daescu A, Daniluc RI, Stoian D. The Paradox of Sexual Dysfunction Observed during Pregnancy. In *Healthcare* 2023 Jul 3 (Vol. 11, No. 13, p. 1914). MDPI.
55. Lee R, Serene T, Tan KH. Guidelines on physical activity & exercise in pregnancy. *Perinatal Society of Singapore*, Singapore. 2020 Jan 10:9-16.
56. Karaahmet AY, Gençtürk N, Boyacıoğlu NE. Does perinatal period pelvic floor muscle exercises affect sexuality and pelvic muscle strength? A systematic review and meta-analysis of randomized controlled trials. *Revista da Associação Médica Brasileira*. 2022 Aug 19;68(8):1033-41.
57. Sobhghol SS, Priddis H, Smith CA, Dahlen HG. The effect of pelvic floor muscle exercise on female sexual function during pregnancy and postpartum: a systematic review. *Sexual Medicine Reviews*. 2019 Jan;7(1):13-28.
58. Cox J, Holden J. *Perinatal mental health: A guide to the Edinburgh Postnatal Depression Scale (EPDS)*. Royal College of Psychiatrists; 2003.
59. Signorello LB, Harlow BL, Chekos AK, Repke JT. Postpartum sexual functioning and its relationship to perineal trauma: a retrospective cohort study of primiparous women. *American journal of obstetrics and gynecology*. 2001 Apr 1;184(5):881-90.
60. Delgado-Pérez E, Rodríguez-Costa I, Vergara-Pérez F, Blanco-Morales M, Torres-Lacombe M. Recovering sexuality after childbirth. What strategies do women adopt? A qualitative study. *International Journal of Environmental Research and Public Health*. 2022 Jan 15;19(2):950.
61. Abdool Z, Thakar R, Sultan AH. Postpartum female sexual function. *European Journal of Obstetrics & Gynecology and Reproductive Biology*. 2009 Aug 1;145(2):133-7.
62. Royal College of Obstetricians and Gynaecologists (RCOG). *The Management of Third- and Fourth-Degree Perineal Tears Green-top Guideline No. 29* [Internet]. 2015. Available from: <https://www.rcog.org.uk/media/5jeb5hzu/gtg-29.pdf>
63. Hady DA, Kassem AR, Abdalla HA. Effect of knack technique on vaginal laxity in multiparous women: A randomized controlled trial. *Physiotherapy Research International*. 2024 Jan;29(1):e2036.

64. Rahmani A, Fallahi A, Allahqoli L, Grylka-Baeschlin S, Alkatout I. How do new mothers describe their postpartum sexual quality of life? a qualitative study. *BMC Women's Health*. 2023 Sep 9;23(1):477.
65. Evenson KR, Brown WJ, Brinson AK, Budzynski-Seymour E, Hayman M. A review of public health guidelines for postpartum physical activity and sedentary behavior from around the world. *Journal of Sport and Health Science*. 2024 Jul 1;13(4):472-83.
66. Mohammed GF, Hashish RK. Sexual violence against females and its impact on their sexual function. *Egyptian Journal of Forensic Sciences*. 2015 Sep 1;5(3):96-102.
67. Yehuda R, Lehrner AM, Rosenbaum TY. PTSD and sexual dysfunction in men and women. *The journal of sexual medicine*. 2015 May;12(5):1107-19.
68. Taylor NM, King CK. Sexual health and the LGBTQ+ community. *Primary Care: Clinics in Office Practice*. 2021 Jun 1;48(2):271-82.
69. Sousa Rodrigues Guedes T, Barbosa Otoni Gonçalves Guedes M, de Castro Santana R, Costa da Silva JF, Almeida Gomes Dantas A, Ochandorena-Acha M, Terradas-Monllor M, Jerez-Roig J, Bezerra de Souza DL. Sexual dysfunction in women with cancer: a systematic review of longitudinal studies. *International Journal of Environmental Research and Public Health*. 2022 Sep 21;19(19):11921.

Note to HCPs: Stay updated with the latest research and recommendations in sexual health to provide evidence-based care. Regular training and professional development opportunities can enhance your skills and comfort in addressing sexual health concerns. Establishing a supportive network among colleagues can facilitate shared learning and improve patient outcomes.

Disclaimer: These guidelines are intended to support, not replace, clinical judgment. Individual patient circumstances may require tailored approaches. Always consider the patient's preferences, cultural background, and specific health needs when applying these guidelines.

The members of the Sexual Health Guidelines for Women of Reproductive Age in Singapore Work Group are:

Chairpersons: Dr Jean-Jasmin Lee Mi-Li, Family Medicine Service, KKH & Dr Tan Tse Yeun, Department of Reproductive Medicine, KKH
 Ms Caroline Chua, Physiotherapy Department, KKH
 Ms Cheong Xinyi, Psychology Service, KKH
 A/Prof Tan Lay Kok, College of Obstetricians and Gynaecologists Singapore
 Dr Jason Lim, Obstetrical and Gynaecological Society of Singapore
 Dr Suraj Kumar, College of Family Physicians Singapore
 Dr Chie Zhi Ying, Singapore Medical Association
 Ms Jay Lim, Psychology Service, KKH
 Dr Chong Weiliang, Singapore Urological Association
 Ms Yip Wan Hui, Singapore Physiotherapy Association
 Dr Melissa Chiang, Health Promotion Board
 Ms Delphine Koh, SingHealth Psychology Committee of Practice
 Dr Huang Zhong Wei, Society for the Study of Andrology and Sexology, Singapore
 Ms June Eng, Physiotherapy Department, KKH

Dr Ku Chee Wai, Department of Reproductive Medicine, KKH

Dr Benny Loo Kai Guo, Sport and Exercise Medicine Service, KKH

Dr Quah Phaik Ling, Senior Research Fellow, Division of Obstetrics and Gynaecology, KKH

Prof Tan Kok Hian, Perinatal Society of Singapore, and Lead IPRAMHO MCHRI, KKH

APPENDIX 1: FEMALE SEXUAL FUNCTION INDEX-6 (FSFI-6). Adapted from Isidori.¹⁸

Over the past 4 weeks:

- | | |
|---|--|
| 1. How would you rate your level (degree) of sexual desire or interest? | 5 = Very high
4 = High
3 = Moderate
2 = Low
1 = Very low or none at all |
| 2. How would you rate your level of sexual arousal ("turn on") during sexual activity or intercourse? | 0 = No sexual activity
5 = Very high
4 = High
3 = Moderate
2 = Low
1 = Very low or none at all |
| 3. How often did you become lubricated ("wet") during sexual activity or intercourse? | 0 = No sexual activity
5 = Almost always or always
4 = Most times
3 = Sometimes
2 = A few times
1 = Almost never or never |
| 4. When you had sexual stimulation or intercourse, how often did you reach orgasm? | 0 = No sexual activity
5 = Almost always or always
4 = Most times
3 = Sometimes
2 = A few times
1 = Almost never or never |
| 5. How satisfied have you been with your overall sexual life? | 5 = Very satisfied
4 = Moderately satisfied
3 = About equally satisfied and dissatisfied
2 = Moderately dissatisfied
1 = Very dissatisfied |
| 6. How often did you experience discomfort or pain during vaginal penetration? | 0 = Did not attempt intercourse
5 = Almost never or never
4 = A few times
3 = Sometimes
2 = Most times
1 = Almost always or always |

Total score:

Note: Total score is the sum of the ordinal responses to the 6 items; the score can range from 2-30 with lower scores corresponding to worse sexual functioning. A cut off score of 19 is indicative of high probability of sexual dysfunction.

Appendix 2 PLSSIT Model

Table 2. PLISSIT Model.¹⁹

The First Level of Treatment
Permission
Encourage patients to express sexual concerns
The Second Level of Treatment
Limited Information
Educate on how changes (e.g., menstrual cycle) can affect sexuality
The Third Level of Treatment
Specific Suggestions
Recommend practical solutions (e.g., lubricants, new intimacy methods)
The Fourth Level of Treatment
Intensive Therapy
Refer for specialized treatment if necessary